



EP 05998



PCT/EP 00 / 05998

27 JUL 2000	
WIPO	PCT INVESTOR IN PEOPLE

10/018695

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

4
4/1

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed *AmBrewer*

Dated 8 June 2000

THIS PAGE BLANK (USPTO)

For Official use

31 JAN 2000

01FEB00 E509492-4 D02029
P01/7700 0.00-0002200.4

Your Reference: FB/RH/ B45209

0002200.4

Notes

Please type, or write in dark ink using CAPITAL letters. A prescribed fee is payable for a request for grant of a patent. For details, please contact the Patent Office (telephone 071-483 4700).

Rule 16 of the Patents Rules 1990 is the main rule governing the completion and filing of this form.

- ② Do not give trading styles, for example, 'Trading as XYZ company', nationality or former names, for example, 'formerly (known as) ABC Ltd' as these are not required.

Warning

After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977 and will inform the applicant if such prohibition or restriction is necessary. Applicants resident in the United Kingdom are also reminded that under Section 23, applications may not be filed abroad without written permission unless an application has been filed not less than 6 weeks previously in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction revoked.

The Patent Office

Request for grant of a

Patent

Form 1/77

Patents Act 1977

① Title of invention

1 Please give the title
of the invention **NOVEL USE**

② Applicant's details

☐ First or only applicant

2a If you are applying as a corporate body please give:
Corporate Name **SmithKline Beecham Biologicals s.a.**

Country (and State
of incorporation, if
appropriate)

Belgium**S781117001**

2b If you are applying as an individual or one of a partnership
please give in full:

Surname

Forenames

2c In all cases, please give the following details:

Address: **Rue de l'Institut 89, B-1330 Rixensart,
Belgium**

UK postcode
(if applicable)

Country **Belgium**
ADP number **5800974002**
(if known)

2d, 2e and 2f: If there are further applicants please provide details on a separate sheet of paper

2d

Second applicant (if any)

If you are applying as a corporate body please give:
Corporate Name

Country (and State of
Incorporation, if appropriate)

2e

If you are applying as an individual or one of a partnership please
give in full:

Surname:

Forenames:

2f

In all cases, please give the following details:

Address:

UK postcode
(if applicable)
Country
ADP number
(if known)

3

An address for
service in the United Kingdom
must be supplied

Please mark correct box.

3

Address for service details

3a

Have you appointed an agent to deal with your application?

Yes ☒

No ☐ go to 3b



please give details below

Agent's name Kathryn L Privett

Agent's address SmithKline Beecham
Corporate Intellectual Property
Two New Horizons Court
Brentford, Middlesex
TW8 9EP

Postcode
Agent's ADP
number

4471231005

3b: If you have appointed an agent,
all correspondence concerning
your application will be sent to the
agent's United Kingdom address.

3b

If you have not appointed an agent please give a name and address
in the United Kingdom to which all correspondence will be sent:

Name

Address

Postcode
number
ADP number
(if known)

Daytime telephone
(if available)


3. Reference number

4. Agent's or
applicant's reference number (if applicable) FB/RH B45029

5 Claiming an earlier application date

5. Are you claiming that this application be treated as having been filed on the date of filing of an earlier application?

Yes ☐

No ☒  go to 6



please give details below

☐ number of earlier
application or patent
number

☐ filing date (day month year)

☐ and the Section of the Patents Act 1977 under which you are
claiming:

15(4) (Divisional) ☐ 8(3) ☐ 12(6) ☐ 37(4) ☐

6 Declaration of priority

6. If you are declaring priority from previous application(s), please give:

Country of Filing

Priority application
number
(if known)

Filing Date
(day, month, year)

Please mark correct box

Please mark correct box

6 If you are declaring priority from a
PCT Application please enter 'PCT'
as the country and enter the country
code (for example, GB) as part of
the application number

*Please give the date in all number
format, for example, 31/05/90 for
31 May 1990*

7

- any applicant is not an inventor
- there is an inventor who is not
- an applicant, or
- any applicant is a corporate body.

- 8 Please supply duplicates of claim(s), abstract, description and drawings).

Please mark correct box(es)

- 9 You or your appointed agent (see Rule 90 of the Patents Rules 1990) must sign this request.

Please sign here ↑

A completed fee sheet should preferably accompany the fee.

7 Inventorship

7. Are you (the applicant or applicants) the sole inventor or the joint inventors?

Please mark correct box

Yes ☐ No ☒

A Statement of Inventorship on Patents form 7/77 will need to be filed (see Rule 15).).

8 Checklist

8a Please fill in the number of sheets for each of the following types of document contained in this application

Continuation sheets for this Patents Form 1/77

Claim(s)

2

Description

42

Abstract

Drawing(s)

21

only

8b Which of the following documents also accompanies the application?

Priority documents (please state how many)

Translation(s) of Priority documents (please state how many)

Patents Form 7/77 - Statement of Inventorship and Right to Grant

Patents Form 9/77 - Preliminary Examination Report

Patents Form 10/77 - Request for Substantive Examination

9 Request

I/We request the grant of a patent on the basis of this application.

Signed

Kathryn L Privett
Kathryn L Privett

Date:

31/01/00

(day)

month

year)

Chartered Patent Attorney
Attorney for the Applicant

Please return the completed form, attachments and duplicates where requested, together with the prescribed fee to either;

☐ The Comptroller or
The Patent Office
Cardiff Road
Newport
Gwent
NP9 1RH

☐ The Comptroller
The Patent Office
25 Southampton Buildings
London
WC2A 1AY

NOVEL USE

DESCRIPTION

The present invention relates to novel uses of HIV proteins in medicine and vaccine compositions containing such HIV proteins. In particular, the invention relates to the use of HIV Tat and HIV gp120 proteins in combination. Furthermore, the invention relates to the use of HIV Nef and HIV gp120 proteins in combination.

HIV-1 is the primary cause of the acquired immune deficiency syndrome (AIDS) which is regarded as one of the world's major health problems. Although extensive research throughout the world has been conducted to produce a vaccine, such efforts thus far have not been successful.

The HIV envelope glycoprotein gp120 is the viral protein that is used for attachment to the host cell. This attachment is mediated by the binding to two surface molecules of helper T cells and macrophages, known as CD4 and one of the two chemokine receptors CCR-4 or CXCR-5. The gp120 protein is first expressed as a larger precursor molecule (gp160), which is then cleaved post-translationally to yield gp120 and gp41. The gp120 protein is retained on the surface of the virion by linkage to the gp41 molecule, which is inserted into the viral membrane.

The gp120 protein is the principal target of neutralizing antibodies, but unfortunately the most immunogenic regions of the proteins (V3 loop) are also the most variable parts of the protein. Therefore, the use of gp120 (or its precursor gp160) as a vaccine antigen to elicit neutralizing antibodies is thought to be of limited use for a broadly protective vaccine. The gp120 protein does also contain epitopes that are recognized by cytotoxic T lymphocytes (CTL). These effector cells are able to eliminate virus-infected cells, and therefore constitute a second major antiviral immune mechanism. In contrast to the target regions of neutralizing antibodies some CTL epitopes appear to be relatively conserved among different HIV strains. For this reason gp120 and gp160 are considered to be useful antigenic components in vaccines that aim at eliciting cell-mediated immune responses (particularly CTL).

Non-envelope proteins of HIV-1 have been described and include for example internal structural proteins such as the products of the *gag* and *pol* genes and, other non-structural proteins such as Rev, Nef, Vif and Tat (Greene et al., New England J. Med, **324**, 5, 308 et seq (1991) and Bryant et al. (Ed. Pizzo), *Pediatr. Infect. Dis. J.*, **11**, 5, 390 et seq (1992).

HIV Tat and Nef proteins are early proteins, that is, they are expressed early in infection and in the absence of structural protein.

In a conference presentation (C. David Pauza, Immunization with Tat toxoid attenuates SHIV89.6PD infection in rhesus macaques, 12th Cent Gardes meeting, Marnes-La-Coquette, 26.10.1999), experiments were described in which rhesus macaques were immunised with Tat toxoid alone or in combination with an envelope glycoprotein gp160 vaccine combination (one dose recombinant vaccinia virus and one dose recombinant protein). However, the results observed showed that the presence of the envelope glycoprotein gave no advantage over experiments performed with Tat alone.

However, we have found that a Tat- and/or Nef-containing immunogen acts synergistically with gp120 in protecting rhesus monkeys from a pathogenic challenge with chimeric human-simian immunodeficiency virus (SHIV). To date the SHIV infection of rhesus macaques is considered to be the most relevant animal model for human AIDS. Therefore, we have used this preclinical model to evaluate the protective efficacy of vaccines containing a gp120 antigen and a Nef- and Tat-containing antigen either alone or in combination. Analysis of two markers of viral infection and pathogenicity, the percentage of CD4-positive cells in the peripheral blood and the concentration of free SHIV RNA genomes in the plasma of the monkeys, indicated that the two antigens acted in synergy. Immunization with either gp120 or NefTat + SIV Nef alone did not result in any difference compared to immunization with an adjuvant alone. In contrast, the administration of the combination of both antigens resulted in a marked improvement of the two above-mentioned parameters in all animals of those particular experimental group.

Thus, according to the present invention there is provided a new use of HIV Tat and/or Nef protein together with HIV gp120 in the manufacture of a vaccine for the prophylactic or therapeutic immunisation of humans against HIV.

As described above, the NefTat protein, the SIV Nef protein and gp120 protein together give an enhanced response over that which is observed when either NefTat + SIV Nef, or gp120 are used alone. This enhanced response, or synergy can be seen in a decrease in viral load as a result of vaccination with these combined proteins. Alternatively, or additionally the enhanced response manifests itself by a maintenance of CD4+ levels over those levels found in the absence of vaccination with HIV

NefTat, SIV Nef and HIV gp120. The synergistic effect is attributed to the combination of gp120 and Tat, or gp120 and Nef, or gp120 and both Nef and Tat.

The HIV Tat protein in the vaccine of the present invention may, optionally be linked to an HIV Nef protein, for example as a fusion protein.

The HIV Tat protein, the HIV Nef protein or the NefTat fusion protein in the present invention may have a C terminal Histidine tail which preferably comprises between 5-10 Histidine residues. The presence of an histidine (or 'His') tail aids purification.

In a preferred embodiment the proteins are expressed with a Histidine tail comprising between 5 to 10 and preferably six Histidine residues. These are advantageous in aiding purification. Separate expression, in yeast (*Saccharomyces cerevisiae*), of Nef (Macreadie I.G. et al., 1993, *Yeast* 9 (6) 565-573) and Tat (Braddock M et al., 1989, *Cell* 58 (2) 269-79) has been reported. Nef protein only is myristilated. The expression of Nef and Tat separately in a *Pichia* expression system (Nef-His and Tat-His constructs), and the expression of a fusion construct Nef-Tat-His have been described previously in WO99/16884.

The DNA and amino acid sequences of representative Nef-His (Seq. ID. No.s 8 and 9), Tat-His (Seq. ID. No.s 10 and 11) and of Nef-Tat-His fusion proteins (Seq. ID. No.s 12 and 13) are set forth in Figure 2.

The HIV proteins of the present invention may be used in their native conformation, or more preferably, may be modified for vaccine use. These modifications may either be required for technical reasons relating to the method of purification, or they may be used to biologically inactivate one or several functional properties of the Tat or Nef protein. Thus the invention encompasses derivatives of HIV proteins which may be, for example mutated proteins. The term 'mutated' is used herein to mean a molecule which has undergone deletion, addition or substitution of one or more amino acids using well known techniques for site directed mutagenesis or any other conventional method.

For example, a mutant Tat protein may be mutated so that it is biologically inactive whilst still maintaining its immunogenic epitopes. One possible mutated tat gene, constructed by D.Clements (Tulane University), (originating from BH10 molecular clone) bears mutations in the active site region (Lys41→Ala) and in RGD motif (Arg78→Lys and Asp80→Glu) (Virology 235: 48-64, 1997).

A mutated Tat is illustrated in Figure 2 (Seq. ID. No.s 22 and 23) as is a Nef-Tat Mutant-His (Seq. ID. No.s 24 and 25).

The HIV Tat or Nef proteins in the vaccine of the present invention may be modified by a chemical method during the purification process to render the proteins stable and monomeric. One method to prevent oxidation of a protein such as Tat or Nef is a chemical modification of the protein's thiol groups after a reduction step known as carboxymethylation. Such chemical modification does not modify functional properties of Tat or Nef as assessed by cell binding assays and inhibition of lymphoproliferation of human peripheral blood mononuclear cells.

The HIV Tat protein and HIV gp120 proteins can be purified by the methods outlined in the attached examples.

The vaccine of the present invention will contain an immunoprotective or immunotherapeutic quantity of the Tat and/or Nef or NefTat and gp120 antigens and may be prepared by conventional techniques.

Vaccine preparation is generally described in New Trends and Developments in Vaccines, edited by Voller et al., University Park Press, Baltimore, Maryland, U.S.A. 1978. Encapsulation within liposomes is described, for example, by Fullerton, U.S. Patent 4,235,877. Conjugation of proteins to macromolecules is disclosed, for example, by Likhite, U.S. Patent 4,372,945 and by Armor et al., U.S. Patent 4,474,757.

The amount of protein in the vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical

vaccinees. Such amount will vary depending upon which specific immunogen is employed. Generally, it is expected that each dose will comprise 1-1000 µg of each protein, preferably 2-200 µg, most preferably 4-40 µg of Tat or Nef or NefTat and most preferably 20-150 µg of gp120. An optimal amount for a particular vaccine can be ascertained by standard studies involving observation of antibody titres and other responses in subjects. Following an initial vaccination, subjects may receive a boost in about 4 weeks.

The proteins of the present invention are preferably adjuvanted in the vaccine formulation of the invention. Adjuvants are described in general in Vaccine Design – the Subunit and Adjuvant Approach, edited by Powell and Newman, Plenum Press, New York, 1995.

Suitable adjuvants include an aluminium salt such as aluminium hydroxide gel (alum) or aluminium phosphate, but may also be a salt of calcium, iron or zinc, or may be an insoluble suspension of acylated tyrosine, or acylated sugars, cationically or anionically derivatised polysaccharides, or polyphosphazenes.

In the formulation of the invention it is preferred that the adjuvant composition induces a preferential Th1 response. However it will be understood that other responses, including other humoral responses, are not excluded.

Preferred Th1-type immunostimulants which may be formulated to form adjuvants suitable for use in the present invention include and are not restricted to the following.

Monophosphoryl lipid A, in particular 3-de-O-acylated monophosphoryl lipid A (3D-MPL), is a preferred Th1-type immunostimulant for use in the invention. 3D-MPL is a well known adjuvant manufactured by Ribi Immunochem, Montana. Chemically it is often supplied as a mixture of 3-de-O-acylated monophosphoryl lipid A with either 4, 5, or 6 acylated chains. It can be prepared by the methods taught in GB 2122204 B. A preferred form of 3D-MPL is in the form of a particulate formulation having a small particle size less than 0.2µm in diameter, and its method of manufacture is disclosed in EP 0 689 454.

Saponins are also preferred Th1 immunostimulants in accordance with the invention. Saponins are well known adjuvants and are taught in: Lacaille-Dubois, M and Wagner H. (1996. A review of the biological and pharmacological activities of saponins. *Phytomedicine* vol 2 pp 363-386). For example, Quil A (derived from the bark of the South American tree *Quillaja Saponaria* Molina), and fractions thereof, are described in US 5,057,540 and "Saponins as vaccine adjuvants", Kensil, C. R., *Crit Rev Ther Drug Carrier Syst*, 1996, 12 (1-2):1-55; and EP 0 362 279 B1. The haemolytic saponins QS21 and QS17 (HPLC purified fractions of Quil A) have been described as potent systemic adjuvants, and the method of their production is disclosed in US Patent No. 5,057,540 and EP 0 362 279 B1. Also described in these references is the use of QS7 (a non-haemolytic fraction of Quil-A) which acts as a potent adjuvant for systemic vaccines. Use of QS21 is further described in Kensil *et al.* (1991. *J. Immunology* vol 146, 431-437). Combinations of QS21 and polysorbate or cyclodextrin are also known (WO 99/10008). Particulate adjuvant systems comprising fractions of QuilA, such as QS21 and QS7 are described in WO 96/33739 and WO 96/11711.

Another preferred immunostimulant is an immunostimulatory oligonucleotide containing unmethylated CpG dinucleotides ("CpG"). CpG is an abbreviation for cytosine-guanosine dinucleotide motifs present in DNA. CpG is known in the art as being an adjuvant when administered by both systemic and mucosal routes (WO 96/02555, EP 468520, Davis *et al.*, *J.Immunol*, 1998, 160(2):870-876; McCluskie and Davis, *J.Immunol.*, 1998, 161(9):4463-6). Historically, it was observed that the DNA fraction of BCG could exert an anti-tumour effect. In further studies, synthetic oligonucleotides derived from BCG gene sequences were shown to be capable of inducing immunostimulatory effects (both in vitro and in vivo). The authors of these studies concluded that certain palindromic sequences, including a central CG motif, carried this activity. The central role of the CG motif in immunostimulation was later elucidated in a publication by Krieg, *Nature* 374, p546 1995. Detailed analysis has shown that the CG motif has to be in a certain sequence context, and that such sequences are common in bacterial DNA but are rare in vertebrate DNA. The immunostimulatory sequence is often: Purine, Purine, C, G, pyrimidine, pyrimidine;

wherein the CG motif is not methylated, but other unmethylated CpG sequences are known to be immunostimulatory and may be used in the present invention.

In certain combinations of the six nucleotides a palindromic sequence is present. Several of these motifs, either as repeats of one motif or a combination of different motifs, can be present in the same oligonucleotide. The presence of one or more of these immunostimulatory sequences containing oligonucleotides can activate various immune subsets, including natural killer cells (which produce interferon γ and have cytolytic activity) and macrophages (Wooldrige et al Vol 89 (no. 8), 1977). Other unmethylated CpG containing sequences not having this consensus sequence have also now been shown to be immunomodulatory.

CpG when formulated into vaccines, is generally administered in free solution together with free antigen (WO 96/02555; McCluskie and Davis, *supra*) or covalently conjugated to an antigen (WO 98/16247), or formulated with a carrier such as aluminium hydroxide ((Hepatitis surface antigen) Davis *et al. supra* ; Brazolot-Millan *et al.*, *Proc.Natl.Acad.Sci.*, USA, 1998, 95(26), 15553-8).

Such immunostimulants as described above may be formulated together with carriers, such as for example liposomes, oil in water emulsions, and or metallic salts, including aluminium salts (such as aluminium hydroxide). For example, 3D-MPL may be formulated with aluminium hydroxide (EP 0 689 454) or oil in water emulsions (WO 95/17210); QS21 may be advantageously formulated with cholesterol containing liposomes (WO 96/33739), oil in water emulsions (WO 95/17210) or alum (WO 98/15287); CpG may be formulated with alum (Davis *et al. supra* ; Brazolot-Millan *supra*) or with other cationic carriers.

Combinations of immunostimulants are also preferred, in particular a combination of a monophosphoryl lipid A and a saponin derivative (WO 94/00153; WO 95/17210; WO 96/33739; WO 98/56414; WO 99/12565; WO 99/11241), more particularly the combination of QS21 and 3D-MPL as disclosed in WO 94/00153. Alternatively, a combination of CpG plus a saponin such as QS21 also forms a potent adjuvant for use in the present invention.

Thus, suitable adjuvant systems include, for example, a combination of monophosphoryl lipid A, preferably 3D-MPL, together with an aluminium salt. An enhanced system involves the combination of a monophosphoryl lipid A and a saponin derivative particularly the combination of QS21 and 3D-MPL as disclosed in WO 94/00153, or a less reactogenic composition where the QS21 is quenched in cholesterol containing liposomes (DQ) as disclosed in WO 96/33739.

A particularly potent adjuvant formulation involving QS21, 3D-MPL & tocopherol in an oil in water emulsion is described in WO 95/17210 and is another preferred formulation for use in the invention.

Another preferred formulation comprises a CpG oligonucleotide alone or together with an aluminium salt.

In another aspect of the invention, the vaccine may contain DNA encoding one or more of the Tat, Nef and gp120 polypeptides, such that the polypeptide is generated *in situ*. The DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Numerous gene delivery techniques are well known in the art, such as those described by Rolland, Crit. Rev. Therap. Drug Carrier Systems 15:143-198, 1998 and references cited therein. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter and terminating signal). When the expression system is a recombinant live microorganism, such as a virus or bacterium, the gene of interest can be inserted into the genome of a live recombinant virus or bacterium. Inoculation and *in vivo* infection with this live vector will lead to *in vivo* expression of the antigen and induction of immune responses. Viruses and bacteria used for this purpose are for instance: poxviruses (e.g; vaccinia, fowlpox, canarypox), alphaviruses (Sindbis virus, Semliki Forest Virus, Venezuelan Equine Encephalitis Virus), adenoviruses, adeno-associated virus, picornaviruses (poliovirus, rhinovirus), herpesviruses (varicella zoster virus, etc), Listeria, Salmonella, Shigella, Neisseria, BCG. These viruses and bacteria can be virulent, or attenuated in various ways in order to obtain live vaccines. Such live vaccines also form part of the invention.

In a further aspect of the present invention there is provided a method of manufacture of a vaccine formulation as herein described, wherein the method comprises mixing a combination of proteins according to the invention. The protein composition may be mixed with a suitable adjuvant and, optionally, a carrier.

Particularly preferred adjuvant and/or carrier combinations for use in the formulations according to the invention are as follows:

- i) 3D-MPL + QS21 in DQ
- ii) Alum + 3D-MPL
- iii) Alum + QS21 in DQ + 3D-MPL
- iv) Alum + CpG
- v) 3D-MPL + QS21 in DQ + oil in water emulsion
- vi) CpG

The invention is illustrated in the accompanying examples and Figures:

EXAMPLES

General

The Nef gene from the Bru/Lai isolate (Cell 40: 9-17, 1985) was selected for the constructs of these experiments since this gene is among those that are most closely related to the consensus Nef .

The starting material for the Bru/Lai Nef gene was a 1170bp DNA fragment cloned on the mammalian expression vector pcDNA3 (pcDNA3/Nef).

The Tat gene originates from the BH10 molecular clone. This gene was received as an HTLV III cDNA clone named pCV1 and described in Science, 229, p69-73, 1985.

The expression of the Nef and Tat genes could be in *Pichia* or any other host.

Example 1. EXPRESSION OF HIV-1 *nef* AND *tat* SEQUENCES IN *PICHIA PASTORIS*.

Nef protein, Tat protein and the fusion Nef -Tat were expressed in the methylotrophic yeast *Pichia pastoris* under the control of the inducible alcohol oxidase (AOX1) promoter.

To express these HIV-1 genes a modified version of the integrative vector PHIL-D2 (INVITROGEN) was used. This vector was modified in such a way that expression of heterologous protein starts immediately after the native ATG codon of the AOX1 gene and will produce recombinant protein with a tail of one glycine and six histidines residues . This PHIL-D2-MOD vector was constructed by cloning an oligonucleotide linker between the adjacent *Asu*II and *Eco*RI sites of PHIL-D2 vector (see Figure 2). In addition to the His tail, this linker carries *Nco*I, *Spe*I and *Xba*I restriction sites between which *nef*, *tat* and *nef-tat* fusion were inserted.

1.1 CONSTRUCTION OF THE INTEGRATIVE VECTORS pRIT14597 (encoding Nef-His protein), pRIT14598 (encoding Tat-His protein) and pRIT14599 (encoding fusion Nef-Tat-His).

The *nef* gene was amplified by PCR from the pcDNA3/Nef plasmid with primers 01 and 02.

NcoI

PRIMER 01 (Seq ID NO 1): 5'ATCGTCCATG.GGT.GGC.AAG.TGG.T 3'

SpeI

PRIMER 02 (Seq ID NO 2): 5' CGGCTACTAGTGCAGTTCTTGAA 3'

The PCR fragment obtained and the integrative PHIL-D2-MOD vector were both restricted by NcoI and SpeI, purified on agarose gel and ligated to create the integrative plasmid pRIT14597 (see Figure 2).

The *tat* gene was amplified by PCR from a derivative of the pCV1 plasmid with primers 05 and 04:

SpeI

PRIMER 04 (Seq ID NO 4): 5' CGGCTACTAGTTTCCTTCGGGCCT 3'

NcoI

PRIMER 05 (Seq ID NO 5): 5'ATCGTCCATGGAGCCAGTAGATC 3'

An NcoI restriction site was introduced at the 5' end of the PCR fragment while a SpeI site was introduced at the 3' end with primer 04. The PCR fragment obtained

and the PHIL-D2-MOD vector were both restricted by NcoI and SpeI, purified on agarose gel and ligated to create the integrative plasmid pRIT14598.

To construct pRIT14599, a 910bp DNA fragment corresponding to the *nef-tat*-His coding sequence was ligated between the EcoRI blunted(T4 polymerase) and NcoI sites of the PHIL-D2-MOD vector. The *nef-tat*-His coding fragment was obtained by XbaI blunted(T4 polymerase) and NcoI digestions of pRIT14596.

1.2 TRANSFORMATION OF PICHIA PASTORIS STRAIN GS115(his4).

To obtain *Pichia pastoris* strains expressing Nef-His, Tat-His and the fusion Nef-Tat-His, strain GS115 was transformed with linear NotI fragments carrying the respective expression cassettes plus the HIS4 gene to complement his4 in the host genome. Transformation of GS115 with NotI-linear fragments favors recombination at the AOXI locus.

Multicopy integrant clones were selected by quantitative dot blot analysis and the type of integration, insertion (Mut⁺ phenotype) or transplacement (Mut^s phenotype), was determined.

From each transformation, one transformant showing a high production level for the recombinant protein was selected :

Strain Y1738 (Mut⁺ phenotype) producing the recombinant Nef-His protein, a myristylated 215 amino acids protein which is composed of:

- °Myristic acid
- °A methionine, created by the use of NcoI cloning site of PHIL-D2-MOD vector
- °205 a.a. of Nef protein(starting at a.a.2 and extending to a.a.206)
- °A threonine and a serine created by the cloning procedure (cloning at SpeI site of PHIL-D2-MOD vector.
- °One glycine and six histidines.

Strain Y1739 (Mut⁺ phenotype) producing the Tat-His protein, a 95 amino acid protein which is composed of:

- °A methionine created by the use of NcoI cloning site
- °85 a.a. of the Tat protein(starting at a.a.2 and extending to a.a.86)
- °A threonine and a serine introduced by cloning procedure
- °One glycine and six histidines

Strain Y1737(Mut^s phenotype) producing the recombinant Nef-Tat-His fusion protein, a myristylated 302 amino acids protein which is composed of:

- °Myristic acid
- °A methionine; created by the use of NcoI cloning site
- °205a.a. of Nef protein(starting at a.a.2 and extending to a.a.206)
- °A threonine and a serine created by the cloning procedure
- °85a.a. of the Tat protein(starting at a.a.2 and extending to a.a.86)
- °A threonine and a serine introduced by the cloning procedure
- °One glycine and six histidines

Example 2. EXPRESSION OF HIV-1 Tat-MUTANT IN PICHIA PASTORIS

A mutant recombinant Tat protein has also been expressed. The mutant Tat protein must be **biologically inactive** while **maintaining** its **immunogenic epitopes**.

A double mutant *tat* gene, constructed by D.Clements (Tulane University) was selected for these constructs.

This *tat* gene (originates from BH10 molecular clone) bears **mutations** in the **active site region** (Lys41→Ala)and in **RGD motif** (Arg78→Lys and Asp80→Glu) (Virology 235: 48-64, 1997).

The mutant *tat* gene was received as a cDNA fragment subcloned between the EcoRI and HindIII sites within a CMV expression plasmid (pCMVLys41/KGE)

2.1 CONSTRUCTION OF THE INTEGRATIVE VECTORS

pRIT14912(encoding Tat mutant-His protein) and pRIT14913(encoding fusion Nef-Tat mutant-His).

The *tat* mutant gene was amplified by PCR from the pCMVLys41/KGE plasmid with primers 05 and 04 (see section 1.1construction of pRIT14598)

An NcoI restriction site was introduced at the 5' end of the PCR fragment while a SpeI site was introduced at the 3' end with primer 04. The PCR fragment obtained and the PHIL-D2-MOD vector were both restricted by NcoI and SpeI, purified on agarose gel and ligated to create the integrative plasmid pRIT14912

To construct pRIT14913, the *tat* mutant gene was amplified by PCR from the pCMVLys41/KGE plasmid with primers 03 and 04.

SpeI

PRIMER 03 (Seq ID NO 3): 5' ATCGTACTAGT.GAG.CCA.GTA.GAT.C 3'

SpeI

PRIMER 04 (Seq ID NO 4): 5' CGGCTACTAGTTTTCCTTCGGGCCT 3'

The PCR fragment obtained and the plasmid pRIT14597 (expressing Nef-His protein) were both digested by SpeI restriction enzyme, purified on agarose gel and ligated to create the integrative plasmid pRIT14913

2.2 TRANSFORMATION OF PICHIA PASTORIS STRAIN GS115.

Pichia pastoris strains expressing Tat mutant-His protein and the fusion Nef-Tat mutant-His were obtained, by applying integration and recombinant strain selection strategies previously described in section 1.2 .

Two recombinant strains producing Tat mutant-His protein ,a 95 amino-acids protein, were selected: Y1775 (Mut⁺ phenotype) and Y1776(Mut^s phenotype).

One recombinant strain expressing Nef-Tat mutant-His fusion protein, a 302 amino-acids protein was selected: Y1774(Mut⁺ phenotype).

Example 3: FERMENTATION OF PICHIA PASTORIS PRODUCING RECOMBINANT TAT-HIS.

A typical process is described in the table hereafter.

Fermentation includes a growth phase (feeding with a glycerol-based medium according to an appropriate curve) leading to a high cell density culture and an induction phase (feeding with a methanol and a salts/micro-elements solution). During fermentation the growth is followed by taking samples and measuring their absorbance at 620 nm. During the induction phase methanol was added via a pump and its concentration monitored by Gas chromatography (on culture samples) and by on-line gas analysis with a Mass spectrometer. After fermentation the cells were recovered by centrifugation at 5020g during 30' at 2-8°C and the cell paste stored at -20°C. For further work cell paste was thawed, resuspended at an OD (at 620 nm) of 150 in a buffer (Na₂HPO₄ pH7 50 mM, PMSF 5%, Isopropanol 4 mM) and disrupted by 4 passages in a DynoMill (room 0.6L, 3000 rpm, 6L/H, beads diameter of 0.40-0.70 mm).

For evaluation of the expression samples were removed during the induction, disrupted and analyzed by SDS-Page or Western blot. On Coomassie blue stained SDS-gels the recombinant Tat-his was clearly identified as an intense band presenting a maximal intensity after around 72-96H induction.

Thawing of a Working seed vial	
↓	
Solid preculture 30°C, 14-16H	<u>Synthetic medium</u> : YNB + glucose + agar
↓	
Liquid preculture in two 2L erlenmeyer 30°C, 200 rpm	<u>Synthetic medium</u> : 2 x 400 ml YNB + glycerol Stop when OD > 1 (at 620 nm)
↓	
Inoculation of a 20L fermentor	5L initial medium (FSC006AA) 3 ml antifoam SAG471 (from Witco) Set-points: Temperature : 30°C Overpressure: 0.3 barg Air flow: 20 Nl/min Dissolved O ₂ : regulated > 40% pH : regulated at 5 by NH ₄ OH
↓	
Fed-batch fermentation: growth phase Duration around 40H,	Feeding with glycerol-based medium FFB005AA Final OD between 200-500 OD (620 nm)
Fed-batch fermentation: induction phase Duration: up to 97H.	Feeding with methanol and with a salt/micro-elements solution (FSE021AB).
↓	
Centrifugation	5020g /30 min / 2-8°C
↓	
Recover cell paste and store at -20°C	
↓	
Thaw cells and resuspend at OD150 (620 nm) in buffer	<u>Buffer</u> : Na ₂ HPO ₄ pH7 50 mM, PMSF 5%, Isopropanol 4 mM
↓	
Cell disruption in Dyno-mill 4 passages	<u>Dyno-mill</u> : (room 0.6L, 3000 rpm, 6L/H, beads diameter of 0.40-0.70 mm).
↓	
Transfer for extraction/purification	

Media used for fermentation:**Solid preculture: (YNB + glucose + agar)**

Glucose:	10 g/l	Na ₂ MoO ₄ .2H ₂ O:	0.0002 g/l	Acide folique:	0.000064 g/l
KH ₂ PO ₄ :	1 g/l	MnSO ₄ .H ₂ O:	0.0004 g/l	Inositol:	0.064 g/l
MgSO ₄ .7H ₂ O:	0.5 g/l	H ₃ BO ₃ :	0.0005 g/l	Pyridoxine:	0.008 g/l
CaCl ₂ .2H ₂ O:	0.1 g/l	KI:	0.0001 g/l	Thiamine:	0.008 g/l
NaCl:	0.1 g/l	CoCl ₂ .6H ₂ O:	0.00009 g/l	Niacine:	0.000032 g/l
FeCl ₃ .6H ₂ O:	0.0002 g/l	Riboflavine:	0.000016 g/l	Panthoténate Ca:	0.008 g/l
CuSO ₄ .5H ₂ O:	0.00004 g/l	Biotine:	0.000064 g/l	Para-aminobenzoic acid:	0.000016 g/l
ZnSO ₄ .7H ₂ O:	0.0004 g/l	(NH ₄) ₂ SO ₄ :	5 g/l	Agar	18 g/l

Liquid preculture (YNB + glycerol)

Glycerol:	2% (v/v)	Na ₂ MoO ₄ .2H ₂ O:	0.0002 g/l	Acide folique:	0.000064 g/l
KH ₂ PO ₄ :	1 g/l	MnSO ₄ .H ₂ O:	0.0004 g/l	Inositol:	0.064 g/l
MgSO ₄ .7H ₂ O:	0.5 g/l	H ₃ BO ₃ :	0.0005 g/l	Pyridoxine:	0.008 g/l
CaCl ₂ .2H ₂ O:	0.1 g/l	KI:	0.0001 g/l	Thiamine:	0.008 g/l
NaCl:	0.1 g/l	CoCl ₂ .6H ₂ O:	0.00009 g/l	Niacine:	0.000032 g/l
FeCl ₃ .6H ₂ O:	0.0002 g/l	Riboflavine:	0.000016 g/l	Panthoténate Ca:	0.008 g/l
CuSO ₄ .5H ₂ O:	0.00004 g/l	Biotine:	0.000064 g/l	Para-aminobenzoic acid:	0.000016 g/l
ZnSO ₄ .7H ₂ O:	0.0004 g/l	(NH ₄) ₂ SO ₄ :	5 g/l		

Initial fermentor charge: (FSC006AA)

(NH ₄) ₂ SO ₄ :	6.4 g/l		
KH ₂ PO ₄ :	9 g/l	Na ₂ MoO ₄ .2H ₂ O:	2.04 mg/l
MgSO ₄ .7H ₂ O:	4.7 g/l	MnSO ₄ .H ₂ O:	4.08 mg/l
CaCl ₂ .2H ₂ O:	0.94 g/l	H ₃ BO ₃ :	5.1 mg/l
FeCl ₃ .6H ₂ O:	10 mg/l	KI:	1.022 mg/l
HCl:	1.67 ml/l	CoCl ₂ .6H ₂ O:	0.91 mg/l
CuSO ₄ .5H ₂ O:	0.408 mg/l	NaCl:	0.06 g/l
ZnSO ₄ .7H ₂ O:	4.08 mg/l	Biotine:	0.534 mg/l

Feeding solution used for growth phase (FFB005AA)


Glycérol:	38.7 % v/v	Na ₂ MoO ₄ .2H ₂ O:	5.7 mg/l
MgSO ₄ .7H ₂ O:	13 g/l	CuSO ₄ .5H ₂ O:	1.13 mg/l
CaCl ₂ .2H ₂ O:	2.6 g/l	CoCl ₂ .6H ₂ O:	2.5 mg/l
FeCl ₃ .6H ₂ O:	27.8 mg/l	H ₃ BO ₃ :	14.2 mg/l
ZnSO ₄ .7H ₂ O:	11.3 mg/l	Biotine:	1.5 mg/l
MnSO ₄ .H ₂ O:	11.3 mg/l	KI:	2.84 mg/l
KH ₂ PO ₄ :	24.93 g/l	NaCl:	0.167 g/l


Feeding solution of salts and micro-elements used during induction (FSE021AB):


KH ₂ PO ₄ :	45 g/l	Na ₂ MoO ₄ .2H ₂ O:	10.2 mg/l
MgSO ₄ .7H ₂ O:	23.5 g/l	MnSO ₄ .H ₂ O:	20.4 mg/l
CaCl ₂ .2H ₂ O:	4.70 g/l	H ₃ BO ₃ :	25.5 mg/l
NaCl:	0.3 g/l	KI:	5.11 mg/l
HCl:	8.3 ml/l	CoCl ₂ .6H ₂ O:	4.55 mg/l
CuSO ₄ .5H ₂ O:	2.04 mg/l	FeCl ₃ .6H ₂ O:	50.0 mg/l
ZnSO ₄ .7H ₂ O:	20.4 mg/l	Biotine:	2.70 mg/l

Example 4: PURIFICATION OF Nef-Tat-His FUSION PROTEIN (PICHIA PASTORIS)

The purification scheme has been developed from 146g of recombinant *Pichia pastoris* cells (wet weight) or 2L Dyno-mill homogenate OD 55. The chromatographic steps are performed at room temperature. Between steps, Nef-Tat positive fractions are kept overnight in the cold room (+4°C); for longer time, samples are frozen at -20°C.







* ratio: 0,5M Arginin for a protein concentration of 1600µg/ml.

Purity

The level of purity as estimated by SDS-PAGE is shown in Figure 3 by Daiichi Silver Staining and in Figure 4 by Coomassie blue G250.


After Superdex200 step:	> 95%
After dialysis and sterile filtration steps:	> 95%


Recovery

51mg of Nef-Tat-his protein are purified from 146g of recombinant *Pichia pastoris* cells (= 2L of Dyno-mill homogenate OD 55)

Example 5: PURIFICATION OF OXIDIZED NEF-TAT-HIS FUSION PROTEIN IN PICHIA PASTORIS

The purification scheme has been developed from 73 g of recombinant *Pichia pastoris* cells (wet weight) or 1 L Dyno-mill homogenate OD 50. The chromatographic steps are performed at room temperature. Between steps, Nef-Tat positive fractions are kept overnight in the cold room (+4°C); for longer time, samples are frozen at -20°C.





Concentration	up to 0,8 mg/ml
	10kDa Omega membrane(Filtron)
↓	
Dialysis (O/N - 4°C)	Buffer: 10 mM PO ₄ pH 6.8 – 150 mM NaCl – 0,5 M Arginin
↓	
Sterile filtration	Millex GV 0,22µm

→ Level of purity estimated by SDS-PAGE is shown in Figure 6 (Daiichi Silver Staining, Coomassie blue G250, Western blotting):

After dialysis and sterile filtration steps: > 95%

→ Recovery (evaluated by a colorimetric protein assay: DOC TCA BCA)


2,8 mg of oxidized Nef-Tat-his protein are purified from 73 g of recombinant *Pichia pastoris* cells (wet weight) or 1 L of Dyno-mill homogenate OD 50.

Example 6: PURIFICATION OF REDUCED TAT-HIS PROTEIN (PICHIA PASTORIS)

The purification scheme has been developed from 160 g of recombinant *Pichia pastoris* cells (wet weight) or 2L Dyno-mill homogenate OD 66. The chromatographic steps are performed at room temperature. Between steps, Tat positive fractions are kept overnight in the cold room (+4°C) ; for longer time, samples are frozen at -20°C.

160 g of <i>Pichia pastoris</i> cells	
↓	
Homogenization	Buffer: +2 L 50 mM PO ₄ pH 7.0 – 4 mM PMSF final OD:66
↓	
Dyno-mill disruption (4 passes)	
↓	
Centrifugation	JA10 rotor / 9500 rpm / 30 min / room temperature
↓	
Dyno-mill Pellet	
↓	

Wash (1h - 4°C)	<u>Buffer</u> : +2 L 10 mM PO ₄ pH 7.5 – 150 mM NaCl - 1% Empigen
↓	
Centrifugation	JA10 rotor / 9500 rpm / 30 min / room temperature
↓	
Pellet	
↓	
Solubilisation (O/N - 4°C)	<u>Buffer</u> : + 660 ml 10 mM PO ₄ pH 7.5 – 150 mM NaCl - 4.0 M GuHCl
↓	
Centrifugation	JA10 rotor / 9500 rpm / 30 min / room temperature
↓	
Reduction (4H - room temperature - in the dark)	+ 0,2 M 2-mercaptoethanesulfonic acid, sodium salt (powder addition) / pH adjusted to 7.5 (with 1 M NaOH solution) before incubation
↓	
Carboxymethylation (1/2 h - room temperature - in the dark)	+ 0,25 M Iodoacetamid (powder addition) / pH adjusted to 7.5 (with 1 M NaOH solution) before incubation
↓	
Immobilized metal ion affinity chromatography on Ni ⁺⁺ -NTA-Agarose (Qiagen - 60 ml of resin)	<u>Equilibration buffer</u> : 10 mM PO ₄ pH 7.5 – 150 mM NaCl - 4.0 M GuHCl <u>Washing buffer</u> : 1) Equilibration buffer 2) 10 mM PO ₄ pH 7.5 – 150 mM NaCl – 6 M Urea 3) 10 mM PO ₄ pH 7.5 – 150 mM NaCl - 6M Urea - 35 mM Imidazol <u>Elution buffer</u> : 10 mM PO ₄ pH 7.5 – 150 mM NaCl – 6 M Urea - 0,5 M Imidazol
↓	
Dilution	Down to an ionic strength of 12 mS/cm <u>Dilution buffer</u> : 20 mM Borate pH 8.5 – 6 M Urea
↓	
Cation exchange chromatography on SP Sephacrose FF (Pharmacia - 30 ml of resin)	<u>Equilibration buffer</u> : 20 mM Borate pH 8.5 - 150 mM NaCl - 6.0 M Urea <u>Washing buffer</u> : Equilibration buffer <u>Elution buffer</u> : 20 mM Borate pH 8.5 – 400 mM



→ Level of purity estimated by SDS-PAGE as shown in Figure 7(Daiichi Silver Staining, Coomassie blue G250, Western blotting):


After dialysis and sterile filtration steps: > 95%


→ Recovery (evaluated by a colorimetric protein assay: DOC TCA BCA)


48 mg of reduced Tat-his protein are purified from 160 g of recombinant Pichia pastoris cells (wet weight) or 2 L of Dyno-mill homogenate OD 66.

Example 7: Purification of oxidized Tat-his protein (Pichia Pastoris)

The purification scheme has been developed from 74 g of recombinant Pichia pastoris cells (wet weight) or 1L Dyno-mill homogenate OD60. The chromatographic steps are performed at room temperature. Between steps, Tat positive fractions are kept overnight in the cold room (+4°C) ; for longer time, samples are frozen at -20°C.







→ Level of purity estimated by SDS-PAGE as shown in Figure 8 (Daiichi Silver Staining, Coomassie blue G250, Western blotting):


After dialysis and sterile filtration steps: > 95%


→ Recovery (evaluated by a colorimetric protein assay: DOC TCA BCA)

19 mg of oxidized Tat-his protein are purified from 74 g of recombinant Pichia pastoris cells (wet weight) or 1 L of Dyno-mill homogenate OD 60.

Example 8: PURIFICATION OF SIV REDUCED NEF-HIS PROTEIN (PICHIA PASTORIS)

The purification scheme has been developed from 340 g of recombinant Pichia pastoris cells (wet weight) or 4 L Dyno-mill homogenate OD 100. The chromatographic steps are performed at room temperature. Between steps, Nef positive fractions are kept overnight in the cold room (+4°C); for longer time, samples are frozen at -20°C.





Sterile filtration

Millex GV 0,22µm

→ Level of purity estimated by SDS-PAGE as shown in Figure 9 (Daiichi Silver Staining, Coomassie blue G250, Western blotting):


After dialysis and sterile filtration steps: > 95%


→ Recovery (evaluated by a colorimetric protein assay: DOC TCA BCA)

20 mg of SIV reduced Nef-his protein are purified from 340 g of recombinant *Pichia pastoris* cells (wet weight) or 4 L of Dyno-mill homogenate OD 100.

Example 9: PURIFICATION OF HIV REDUCED NEF-HIS PROTEIN (PICHIA PASTORIS)

The purification scheme has been developed from 160 g of recombinant *Pichia pastoris* cells (wet weight) or 3 L Dyno-mill homogenate OD 50. The chromatographic steps are performed at room temperature. Between steps, Nef positive fractions are kept overnight in the cold room (+4°C); for longer time, samples are frozen at -20°C.





→ Level of purity estimated by SDS-PAGE as shown in Figure 10 (Daichi Silver Staining, Coomassie blue G250, Western blotting):

After dialysis and sterile filtration steps: > 95%

→ Recovery (evaluated by a colorimetric protein assay: DOC TCA BCA)

20 mg of HIV reduced Nef-his protein are purified from 160 g of recombinant *Pichia pastoris* cells (wet weight) or 3 L of Dyno-mill homogenate OD 50.

Example 10: EXPRESSION OF SIV *nef* SEQUENCE IN *PICHTIA PASTORIS*

In order to evaluate Nef and Tat antigens in the pathogenic SHIV challenge model, we have expressed the Nef protein of simian immunodeficiency virus (SIV) of macaques, SIVmac239 (Aids Research and Human Retroviruses, 6:1221-1231,1990).

In the Nef coding region, SIV mac 239 has an in-frame stop codon after 92aa predicting a truncated product of only 10kD. The remainder of the Nef reading frame is open and would be predicted to encode a protein of 263aa (30kD) in its fully open form.

Our starting material for SIVmac239 *nef* gene was a DNA fragment corresponding to the complete coding sequence, cloned on the LX5N plasmid (received from Dr R.C. Desrosiers, Southborough,MA,USA).

This SIV *nef* gene is mutated at the premature stop codon (nucleotide G at position 9353 replaces the original T nucleotide) in order to express the full-length SIVmac239 Nef protein.

To express this SIV *nef* gene in *Pichia pastoris*, the PHIL-D2-MOD Vector (previously used for the expression of HIV-1 *nef* and *tat* sequences) was used. The recombinant protein is expressed under the control of the inducible alcohol oxidase (AOX1) promoter and the c-terminus of the protein is elongated by a Histidine affinity tail that will facilitate the purification.

10.1 CONSTRUCTION OF THE INTEGRATIVE VECTOR pRIT 14908

To construct pRIT 14908, the SIV *nef* gene was amplified by PCR from the pLX5N/SIV-NEF plasmid with primers SNEF1 and SNEF2.

PRIMER SNEF1: 5'ATCGTCCATG.GGTGGAGCTATTTT 3'
NcoI

PRIMER SNEF2: 5' CGGCTACTAGTGCGAGTTTCCTT 3'
SpeI

The SIV *nef* DNA region amplified starts at nucleotide 9077 and terminates at nucleotide 9865 (Aids Research and Human Retroviruses, 6:1221-1231,1990).

An NcoI restriction site (with carries the ATG codon of the *nef* gene) was introduced at the 5' end of the PCR fragment while a SpeI site was introduced at the 3' end. The PCR fragment obtained and the integrative PHIL-D2-MOD vector were both restricted by NcoI and SpeI. Since one NcoI restriction site is present on the SIV *nef* amplified sequence (at position 9286), two fragments of respectively ± 200 bp and ± 600 bp were obtained, purified on agarose gel and ligated to PHIL-D2-MOD vector. The resulting recombinant plasmid received, after verification of the *nef* amplified region by automated sequencing, the pRIT 14908 denomination.

10.2 TRANSFORMATION OF PICHIA PASTORIS STRAIN GS115(his4).

To obtain *Pichia pastoris* strain expressing SIV *nef*-His, strain GS115 was transformed with a linear NotI fragment carrying only the expression cassette and the HIS4 gene (Fig.11).

This linear NotI DNA fragment ,with homologies at both ends with AOX1 resident *P.pastoris* gene, favors recombination at the AOX1 locus.

Multicopy integrant clones were selected by quantitative dot blot analysis .

One transformant showing the best production level for the recombinant protein was selected and received the **Y1772** denomination.

Strain **Y1772** produces the recombinant SIV Nef-His protein, a **272 amino acids** protein which would be composed of:

°Myristic acid

°A methionine, created by the use of NcoI cloning site of PHIL-D2-MOD vector .

°262 amino acids (aa) of Nef protein (starting at aa 2 and extending to aa 263, see Figure 12)

°A threonine and a serine created by the cloning procedure (cloning at SpeI site of PHIL-D2-MOD vector (Fig.11).

°One glycine and six histidines.

Nucleic and Protein sequences are shown on figure 12.

10.3 CHARACTERIZATION OF THE EXPRESSED PRODUCT OF STRAIN Y1772.

Expression level

After 16 hours induction in medium containing 1% methanol as carbon source, abundance of the recombinant Nef-His protein, was estimated at 10% of total protein (Fig.13 , lanes 3-4).

Solubility

Induced cultures of recombinant strain Y1772 producing the Nef-His protein were centrifuged. Cell pellets were resuspended in breaking buffer, disrupted with 0.5mm glass beads and the cell extracts were centrifuged. The proteins contained in the insoluble pellet (P) and in the soluble supernatant (S) were compared on a Coomassie Blue stained SDS-PAGE10%.

As shown in figure 13, the majority of the recombinant protein from strain Y1772 (lanes 3-4) is associated with the insoluble fraction.

Strain Y1772 which presents a satisfactory recombinant protein expression level is used for the production and purification of SIV Nef-His protein.

Example 11: EXPRESSION OF GP120 IN CHO

A stable CHO-K1 cell line which produces a recombinant gp120 glycoprotein has been established. Recombinant gp120 glycoprotein is a recombinant truncated form of the gp120 envelope protein of HIV-1 isolate W61D. The protein is excreted into the cell culture medium, from which it is subsequently purified.

Construction of gp120 transfection plasmid pRIT13968

The envelope DNA coding sequence (including the 5'exon of tat and rev) of HIV-1 isolate W61D was obtained (Dr. Tersmette, CCB, Amsterdam) as a genomic gp160

envelope containing plasmid W61D (Nco-XhoI). The plasmid was designated pRIT13965.

In order to construct a gp120 expression cassette a stop codon had to be inserted at the amino acid glu 515 codon of the gp160 encoding sequence in pRIT13965 using a primer oligonucleotide sequence (DIR 131) and PCR technology. Primer DIR 131 contains three stop codons (in all open reading frames) and a SalI restriction site.

The complete gp120 envelope sequence was then reconstituted from the N-terminal BamHI-DraI fragment (170 bp) of a gp160 plasmid subclone pW61d env (pRIT13966) derived from pRIT13965, and the DraI-SalI fragment (510 bp) generated by PCR from pRIT13965. Both fragments were gel purified and ligated together into the E.coli plasmid pUC18, cut first by SalI (klenow treated), and then by BamHI. This resulted in plasmid pRIT13967. The gene sequence of the XmaI-SalI fragment (1580 bp) containing the gp120 coding cassette was sequenced and found to be identical to the predicted sequence. Plasmid RIT13967 was ligated into the CHO GS-expression vector pEE14 (Celltech Ltd., UK) by cutting first with BclI (klenow treated) and then by XmaI. The resulting plasmid was designated pRIT13968.

Preparation of Master Cell Bank

The gp120-construct (pRIT13968) was transfected into CHO cells by the classical CaPO₄-precipitation/glycerol shock procedure. Two days later the CHOK1 cells were subjected to selective growth medium (GMEM + methionine sulfoximine (MSX) 25 µM + Glutamate + asparagine + 10% Foetal calf serum). Three chosen transfectant clones were further amplified in 175m² flasks and few cell vials were stored at -80°C. C-env 23,9 was selected for further expansion.

A small prebank of cells was prepared and 20 ampoules were frozen. For preparation of the prebank and the MCB, cells were grown in GMEM culture medium, supplemented with 7.5 % fetal calf serum and containing 50 µM MSX. These cell cultures were tested for sterility and mycoplasma and proved to be negative.

The Master Cell Bank CHOK1 env 23.9 (at passage 12) was prepared using cells derived from the premaster cell bank. Briefly, two ampoules of the premaster seed were seeded in medium supplemented with 7.5% dialysed foetal bovine serum. The cells were distributed in four culture flasks and cultured at 37°C. After cell attachment the culture medium was changed with fresh medium supplemented with 50 µM MSX. At confluence, cells were collected by trypsinisation and subcultured with a 1/8 split ratio in T-flasks - roller bottle - cell factory units. Cells were collected from cell factory units by trypsinisation and centrifugation. The cell pellet was resuspended in culture medium supplemented with DMSO as cryogenic preservative. Ampoules were prelabelled, autoclaved and heat-sealed (250 vials). They were checked for leaks and stored overnight at -70°C before storage in liquid nitrogen.

Cell Culture And Production Of Crude Harvest

Two vials from a master cell bank are thawed rapidly. Cells are pooled and inoculated in two T-flasks at $37^{\circ} \pm 1^{\circ}\text{C}$ with an appropriate culture medium supplemented with 7.5 % dialysed foetal bovine (FBS) serum. When reaching confluence (passage 13), cells are collected by trypsinisation, pooled and expanded in 10 T-flasks as above. Confluent cells (passage 14) are trypsinised and expanded serially in 2 cell factory units (each 6000 cm²; passage 15), then in 10 cell factories (passage 16). The growth culture medium is supplemented with 7.5 % dialysed foetal bovine (FBS) serum and 1% MSX. When cells reach confluence, the growth culture medium is discarded and replaced by "production medium" containing only 1 % dialysed foetal bovine serum and no MSX. Supernatant is collected every two days (48 hrs-interval) for up to 32 days. The harvested culture fluids are clarified immediately through a 1.2-0.22 µm filter unit and kept at -20°C before purification.

Example 12: PURIFICATION OF HIV GP 120 (W61D CHO) FROM CELL CULTURE FLUID

All purification steps are performed in a cold room at 2-8°C. pH of buffers are adjusted at this temperature and are filtered on 0.2 µm filter. They are tested for pyrogen content (LAL assay). Optical density at 280 nm, pH and conductivity of column eluates are continuously monitored.

(i) Clarified Culture Fluid

The harvested clarified cell culture fluid (CCF) is filter-sterilized and Tris buffer, pH 8.0 is added to 30 mM final concentration. CCF is stored frozen at -20°C until purification.

(ii) Hydrophobic Interaction Chromatography

After thawing, ammonium sulphate is added to the clarified culture fluid up to 1 M. The solution is passed overnight on a TSK/TOYOPEARL-BUTYL 650 M (TOSOHAAS) column, equilibrated in 30 mM Tris buffer- pH 8.0 - 1 M ammonium sulphate. Under these conditions, the antigen binds to the gel matrix. The column is washed with a decreasing stepwise ammonium sulphate gradient. The antigen is eluted at 30 mM Tris buffer- pH 8.0 - 0.25 M ammonium sulphate.

(iii) Anion-exchange Chromatography

After reducing the conductivity of the solution between 5 and 6 mS/cm, the gP120 pool of fractions is loaded onto a Q-sepharose Fast Flow (Pharmacia) column, equilibrated in Tris-saline buffer - pH 8.0. The column is operated on a negative mode, i.e. gP120 does not bind to the gel, while most of the impurities are retained.

(iv) Concentration and diafiltration by ultrafiltration

In order to increase the protein concentration, the gP120 pool is loaded on a FILTRON membrane "Omega Screen Channel", with a 50 kDa cut-off. At the end of the concentration, the buffer is exchanged by diafiltration with 5 mM phosphate buffer containing CaCl_2 0.3 mM, pH 7.0. If further processing is not performed immediately, the gP120 pool is stored frozen at -20°C. After thawing the solution is filtered onto a 0.2 μM membrane in order to remove insoluble material.

(v) Chromatography on hydroxyapatite

The gP120 UF pool is loaded onto a macro-Prep Ceramic Hydroxyapatite, type II (Biorad) column equilibrated in 5 mM phosphate buffer + CaCl_2 0.3 mM, pH 7.0. The column is washed with the same buffer. The antigen passes through the column and impurities bind to the column.

(vi) Cation exchange chromatography

The gP120 pool is loaded on a CM/TOYOPEARL-650 S (TOSHAAS) column equilibrated in acetate buffer 20 mM, pH 5.0. The column is washed with the same buffer, then acetate 20 mM, pH 5.0 and NaCl 10 mM. The antigen is then eluted by the same buffer containing 80 mM NaCl.

(vii) Ultrafiltration

In order to augment the virus clearance capacity of the purification process, an additional ultrafiltration step is carried out. The gP120 pool is subjected to ultrafiltration onto a FILTRON membrane "Omega Screen Channel", cut-off 150 kDa. This pore-size membrane does not retain the antigen. After the process, the diluted antigen is concentrated on the same type of membrane (Filtron) but with a cut-off of 50 kDa.

(viii) Size exclusion Gel Chromatography

The gP120 pool is applied to a SUPERDEX 200 (PHARMACIA) column in order to exchange the buffer and to eliminate residual contaminants. The column is eluted with phosphate buffer saline (PBS).

(ix) Sterile filtration and storage

Fractions are sterilized by filtration on a 0.2 μ M PVDF membrane (Millipore). After sterile filtration, the purified bulk is stored frozen at -20°C up to formulation. The purification scheme is summarized by the flow sheet below.


⇒ Level of purity of the purified bulk estimated by SDS-PAGE analysis (Silver staining / Coomassie Blue / Western Blotting) is $\geq 95\%$.

⇒ Production yield is around 2.5 mg /L CCF (according to Lowry assay) - Global purification yield is around 25% (according to Elisa assay)

⇒ Purified material is stable 1 week at 37°C (according to WB analysis)

Purification of gP120 from culture fluid

Mark ✓ indicate steps that are critical for virus removal.



Example 13: VACCINE PREPARATION

A vaccine prepared in accordance with the invention comprises the expression products of one or more DNA recombinants encoding an antigen. Furthermore, the formulations comprise a mixture of 3 de -O-acylated monophosphoryl lipid A 3D-MPL and QS21 in an oil/water emulsion or an oligonucleotide containing unmethylated CpG dinucleotide motifs and aluminium hydroxide as carrier.

3D-MPL: is a chemically detoxified form of the lipopolysaccharide (LPS) of the Gram-negative bacteria *Salmonella minnesota*.

Experiments performed at Smith Kline Beecham Biologicals have shown that 3D-MPL combined with various vehicles strongly enhances both the humoral immunity and a T_{H1} type of cellular immunity.

QS21: is a saponin purified from a crude extract of the bark of the *Quillaja Saponaria* Molina tree, which has a strong adjuvant activity: it induces both antigen-specific lymphoproliferation and CTLs to several antigens.

Experiments performed at Smith Kline Beecham Biologicals have demonstrated a clear synergistic effect of combinations of 3D-MPL and QS21 in the induction of both humoral and T_{H1} type cellular immune responses.

The oil/water emulsion is composed of 2 oils (a tocopherol and squalene), and of PBS containing Tween 80 as emulsifier. The emulsion comprises 5% squalene, 5% tocopherol, 2% Tween 80 and has an average particle size of 180 nm (see WO 95/17210).

Experiments performed at Smith Kline Beecham Biologicals have proven that the adjunction of this O/W emulsion to 3D-MPL/QS21 further increases their immunostimulant properties.

Preparation of the oil/water emulsion (2 fold concentrate)

Tween 80 is dissolved in phosphate buffered saline (PBS) to give a 2% solution in the PBS. To provide 100ml two fold concentrate emulsion 5g of DL alpha tocopherol and 5ml of squalene are vortexed to mix thoroughly. 90ml of PBS/Tween solution is added and mixed thoroughly. The resulting emulsion is then passed through a syringe and finally microfluidised by using an M110S Microfluidics machine. The resulting oil droplets have a size of approximately 180 nm.

Preparation of oil in water formulation.

Antigens (100 µg gp120, 20 µg NefTat, and 20 µg SIV Nef, alone or in combination) were diluted in 10 fold concentrated PBS pH 6.8 and H₂O before consecutive addition of SB62, 3D-MPL (50µg), QS21 (50µg) and 1 µg/ml thiomersal as preservative at 5 min interval. The emulsion volume is equal to 50% of the total volume (250µl for a dose of 500µl).

All incubations were carried out at room temperature with agitation.

CpG oligonucleotide (CpG) is a synthetic unmethylated oligonucleotide containing one or several CpG sequence motifs. CpG is a very potent inducer of T_{H1} type immunity compared to the oil in water formulation that induces mainly a mixed T_{H1}/T_{H2} response. CpG induces lower level of antibodies than the oil in water formulation and a good cell mediated immune response. CpG is expected to induce lower local reactogenicity.

Preparation of CpG oligonucleotide solution: CpG dry powder is dissolved in H₂O to give a solution of 5 mg/ml CpG.

Preparation of CpG formulation.

The 3 antigens were dialyzed against NaCl 150 mM to eliminate the phosphate ions that inhibit the adsorption of gp120 on aluminium hydroxide.

The antigens diluted in H₂O (100 µg gp120, 20 µg NefTat and 20 µg SIV Nef) were incubated with the CpG solution (500 µg CpG) for 30 min before adsorption on Al(OH)₃ to favor a potential interaction between the His tail of NefTat and Nef antigens and the oligonucleotide (stronger immunostimulatory effect of CpG

described when bound to the antigen compared to free CpG). Then were consecutively added at 5 min interval $\text{Al}(\text{OH})_3$ (500 μg), 10 fold concentrated NaCl and 1 $\mu\text{g}/\text{ml}$ thiomersal as preservative.

All incubations were carried out at room temperature with agitation.

Example 14: IMMUNIZATION AND SHIV CHALLENGE EXPERIMENT IN RHESUS MONKEYS.

Groups of 4 rhesus monkeys were immunized intramuscularly at 0, 1 and 3 months with the following vaccine compositions:

Group 1:	Adjuvant2	+ gp120		
Group 2:	Adjuvant2	+ gp120	+ NefTat	+ SIV Nef
Group 3:	Adjuvant2		+ NefTat*	+ SIV Nef
Group 4	Adjuvant6	+ gp120	+ NefTat	+ SIV Nef
Group 5	Adjuvant2		+ NefTat	+ SIV Nef
Group 6	Adjuvant2			

Adjuvant 2 comprises squalene/tocopherol/Tween 80/3D-MPL/QS21 and Adjuvant 6 comprisesalum and CpG.

One month after the last immunization all animals were challenged with a pathogenic SHIV (strain 89.6p). From the week of challenge (wk16) blood samples were taken periodically at the indicated time points to determine the % of CD4-positive cells among peripheral blood mononuclear cells by FACS analysis (Figure 14) and the concentration of RNA viral genomes in the plasma by bDNA assay (Figure 15)

CLAIMS

1. Use of HIV Tat protein and HIV gp120 in the manufacture of a vaccine for the prophylactic or therapeutic immunisation of humans against HIV.
2. Use of HIV Tat protein and HIV gp120 as claimed in claim 1 wherein Tat and gp120 in the vaccine act in synergy in the treatment or prevention of HIV.
3. Use of HIV Tat protein and HIV gp120 as claimed in claim 1 or claim 2 wherein the vaccine in use reduces the HIV viral load in HIV infected humans.
4. Use of HIV Tat protein and HIV gp120 as claimed in claims 1 or 2 wherein the vaccine in use results in a maintenance of CD4+ levels over those levels found in the absence of vaccination with HIV Tat protein and HIV gp120.
5. Use of HIV Tat protein and HIV gp120 as claimed in any one of claims 1 – 4 wherein the Tat protein is linked to an HIV Nef protein.
6. Use of HIV Tat protein and HIV gp120 as claimed in any one of claims 1 – 4 wherein the Tat protein is carboxymethylated.
7. Use of HIV Tat protein and HIV gp120 as claimed in any one of claims 1 – 4 wherein the Tat protein is a mutated Tat protein.
8. Use of HIV Tat protein and HIV gp120 as claimed in any one of claims 1 – 4 wherein the Tat protein is oxidised.
9. Use of HIV Tat protein and HIV gp120 as claimed in any one of claims 1 – 4 wherein the Tat protein is reduced.
10. Use of HIV Tat protein and HIV gp120 in a vaccine as claimed in any one of claims 1 – 9 which additionally comprises an adjuvant.
11. Use of HIV Tat protein and HIV gp120 in a vaccine as claimed in claim 10 wherein the adjuvant is a TH1 inducing adjuvant.
12. Use of HIV Tat protein and HIV gp120 in a vaccine as claimed in claim 10 or claim 11 wherein the adjuvant comprises monophosphoryl lipid A or a derivative thereof such as 3-de-O-acylated monophosphoryl lipid A.
13. Use of HIV Tat protein and HIV gp120 in the manufacture of a vaccine as claimed in any one of claims 10 – 12 additionally comprising a saponin adjuvant.
14. Use of HIV Tat protein and HIV gp120 in the manufacture of a vaccine as claimed in any one of claims 10 – 13 additionally comprising an oil in water emulsion.
15. Use of HIV Tat protein and HIV gp 120 in the manufacture of a vaccine as claimed in claim 10 or claim 11 wherein the adjuvant comprises CpG.

16. Use of HIV Nef protein and HIV gp120 in the manufacture of a vaccine for the prophylactic or therapeutic immunisation of humans against HIV.

17. A method of immunising a human against HIV by administering to the human a vaccine comprising HIV Tat and/or Nef protein and HIV gp120.

18. A vaccine composition for human use which vaccine composition comprises HIV Tat and/or Nef protein and HIV gp120.

THIS PAGE BLANK (USPTO)

FIGURE 1

The DNA and amino acid sequences of Nef-His; Tat-His; Nef-Tat-His fusion and mutated Tat is illustrated.

Pichia-expressed constructs (plain constructs)

\Rightarrow Nef - HIS

DNA sequence (Seq. ID. No. 8)

ATGGGTGGCAAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGA
ATGAGACGAGCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAA
AAACATGGAGCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGG
CTAGAAGCACAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTA
AGACCAATGACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTTAAAAGAAAAGGGG
GGACTGGAAGGGCTAATTCCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATC
TACCACACACAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTC
AGATATCCACTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAG
GTAGAAGAGGCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCAT
GGAATGGATGACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCA
TTTCATCACGTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGGC
CACCATCACCATCACCATTAA

Protein sequence (Seq. ID. No. 9)

MGGKWSKSSVVGWPTVRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAW
LEAQEEEEVGFVPTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDLWI
YHTQGYFPDWQNYTPGPGVRYPLTFGWICYKLVPEPDKVEEANKGENTSLLHPVSLH
GMDDPEREVLEWRFD SRLAFHHVARELHPEYFKNCTSGHHHHHH.

\Rightarrow Tat - HIS

DNA sequence (Seq. ID. No. 10)

ATGGAGCCAGTAGATCCTAGACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAA
ACTGCTTGTACCAATTGCTATTGTAAAAAGTGTTGCTTTCATTGCCAAGTTTGTTC
ATAACAAAAGCCTTAGGCATCTCCTATGGCAGGAAGAAGCGGAGACAGCGACGAAGA
CCTCCTCAAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAA

THIS PAGE BLANK (USPTO)

TCCCGAGGGGACCCGACAGGCCCGAAGGAACTAGTGGCCACCATCACCATCACCAT
TAA

Protein sequence (Seq. ID. No. 11)

MEPVDPRLPEWKHPGSQPKTACTNICYCKKCCFHCQVCFITKALGISYGRKKRRQRRR
PPQGSQTHQVSLSKQPTSQSRGDPTGPKETSGHHHHHH.

⇒ Nef - Tat - HIS

DNA sequence (Seq. ID. No. 12)

ATGGGTGGCAAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGA
ATGAGACGAGCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAA
AAACATGGAGCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGG
CTAGAAGCACAAAGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTA
AGACCAATGACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTAAAAGAAAAGGGG
GGAAGTGAAGGGCTAATTCACTCCCAACGAAGACAAGATATCCTTGATCTGTGGATC
TACCACACACAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTC
AGATATCCACTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAG
GTAGAAGAGGCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCAT
GGAATGGATGACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCA
TTTCATCACGTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGAG
CCAGTAGATCCTAGACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAAAGTGT
TGTACCAATTGCTATTGTAAAAAGTGTGCTTTCATTGCCAAGTTTGTTCATAACA
AAAGCCTTAGGCATCTCCTATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCT
CAAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCCGA
GGGGACCCGACAGGCCCGAAGGAACTAGTGGCCACCATCACCATCACCATTAA

Protein sequence (Seq. ID. No. 13)

..
MGGKWSKSSVVGWPTVRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAW
LEAQEEEEVGFVPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWI
YHTQGYFPDWQNYTPGPGVRYPLTFGWCYKLVPVEPDKVEEANKGENTSLLHPVSLH
GMDDPEREVLEWRFD SRLAFHHVARELHPEYFKNCTSEPVDPRLEPWKHPGSQPKTA
CTNICYCKKCCFHCQVCFITKALGISYGRKKRRQRRRPPQGSQTHQVSLSKQPTSQSR
GDPTGPKETSGHHHHHH.

E.coli-expressed constructs (fusion constructs)

⇒ LipoD-Nef-HIS

THIS PAGE BLANK (USPTO)

DNA sequence (Seq. ID. No. 14)

Nucleotides corresponding to the Prot D Fusion Partner are in bold.
The Lipidation Signal Sequence is underlined. After processing, the cysteine coded by the TGT codon, indicated with a star, becomes the amino terminal residue which is then modified by covalently bound fatty acids.

ATGGATCCAAAACTTTAGCCCTTTCTTTATTAGCAGCTGGCGTACTAGCAGGTTGT
AGCAGCCATTTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAAATCATTATT
GCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATACGTTAGAATCTAAAGCACTT
GCTTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACTAAGGATGGT
CGTTTAGTGGTTATTCACGATCACTTTTTAGATGGCTTGACTGATGTTGCGAAAAAA
TTCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTTACCTTAAAA
GAAATTCAAAGTTTAGAAATGACAGAAAACCTTTGAAACCATGGGTGGCAAGTGGTCA
AAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGAGCTGAGCCA
GCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGAGCAATCACA
AGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCACAAGAGGAG
GAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATGACTTACAAG
GCAGCTGTAGATCTTAGCCACTTTTTAAAAGAAAAGGGGGGACTGGAAGGGCTAATT
CACTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACACAAGGCTAC
TTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTGAGATATCCACTGACCTTT
GGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAA
GGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGATGACCCTGAG
AGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTTCATCACGTGGCCCGA
GAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGGCCACCATCACCATCACCAT
TAA

Protein sequence of the processed lipidated ProtD-Nef-HIS protein (Seq. ID. No. 15)

(Amino-acids corresponding to Prot D fusion partner are in bold)

CSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYLEQDLAMTKD
GRLVVIHDHFLDGLTDVAKKFPHRHRKDGRIYVIDFTLKEIQSLEMTENFETMGGKW
SKSSVVGWPTVRRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAWLEAQE
EEVGFPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWYHTQG
YFPDWQNYTPGPGVRYPLTFGWCYKLVPVEPDKVEEANKGENTSLLHPVSLHGMDDP
EREVLEWRFD SRLAFHHVARELHPEYFKNCTSGHHHHHH.

⇒ LipoD-Nef-Tat-HIS

DNA sequence (Seq. ID. No. 16)

THIS PAGE BLANK (USPTO)

Nucleotides corresponding to the Prot D Fusion Partner are in bold.
 The Lipidation Signal Sequence is underlined. After processing, the cysteine coded by the TGT codon, indicated with a star, becomes the amino terminal residue which is then modified by covalently bound fatty acids.

ATGGATCCAAAACTTTAGCCCTTTCTTTATTAGCAGCTGGCGTACTAGCAGGTTGT*
 AGCAGCCATTTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAAATCATTATT
 GCTCACCGTGGTGCTAGCGGTTA^{*}TTTACCAGAGCATACTAGTAATCTAAAGCACTT
 GCGTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACTAAGGATGGT
 CGTTTAGTGGTTATTCACGATCACTTTTTAGATGGCTTGACTGATGTTGCGAAAAAA
 TTCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTTACCTTAAAA
 GAAATTCAAAGTTTAGAAATGACAGAAAACCTTTGAAACCATGGGTGGCAAGTGGTCA
 AAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGGAAAGAATGAGACGAGCTGAGCCA
 GCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGAGCAATCACA
 AGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCACAAGAGGAG
 GAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATGACTTACAAG
 GCAGCTGTAGATCTTAGCCACTTTTTTAAAAGAAAAGGGGGGACTGGAAGGGCTAATT
 CACTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACACAAGGCTAC
 TTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTGAGATATCCACTGACCTTT
 GGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAAA
 GGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGATGACCCTGAG
 AGAGAAAGTGTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTCATCACGTGGCCCGA
 GAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGAGCCAGTAGATCCTAGACTA
 GAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAAACCTGCTTGTTACCAATTGCTATTGT
 AAAAAGTGTTGCTTTTCATTGCCAAGTTTGTTCATAACAAAAGCCTTAGGCATCTCC
 TATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCTCAAGGCAGTCAGACTCAT
 CAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCCGAGGGGACCCGACAGGCCCG
 AAGGAACTAGTGGCCACCATCACCATCACCATTAA

Protein sequence of the processed lipidated ProtD-NEF-TAT-HIS protein (Seq. ID. No. 17)

(Amino-acids corresponding to Prot D fusion partner are in bold)

CSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYLEQDLAMTKD
 GRLVVIHDHFLDGLTDVAKKFPHRHRKDGRIYVIDFTLKEIQSLEMTENFETMGGKW
 SKSSVVGWPTVRERMRRRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAWLEAQE
 EEEVGFPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWIIYHTQG
 YFPDWQNYTPGPGVRYPLTFGWICYKLVPEPDKVEEANKGENTSLLHPVSLHGMDDP
 EREVLEWRFD SRLAFHHVARELHPEYFKNCTSEPVDPRLEPWKHPGSQPKTACTNCY
 CKKCCFHCQVCFITKALGISYGRKKRRQRRRPQGSQTHQVSLSKQPTSQSRGDPTG
 PKETSGHHHHHH.

THIS PAGE BLANK (USPTO)

⇒ ProtD-Nef-HIS

DNA sequence (Seq. ID. No. 18)

Nucleotides corresponding to the Prot D Fusion Partner are in bold.

ATGGATCCAAGCAGCCATTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAA
ATCATTATTGCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATACGTTAGAATCT
AAAGCACTTGCGTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACT
AAGGATGGTCGTTTAGTGTTATTCACGATCACTTTTGTAGATGGCTTGACTGATGTT
GCGAAAAAATTCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTT
ACCTTAAAAGAAATTCAAAGTTTAGAAATGACAGAAAACCTTTGAAACCATGGGTGGC
AAGTGGTCAAAAAGTAGTGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGA
GCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGA
GCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCA
CAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATG
ACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTTAAAAGAAAAGGGGGGACTGGAA
GGGCTAATTCACCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACA
CAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTCAGATATCCA
CTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAG
GCCAATAAAGGAGAGAAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGCAT
GACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTTCATCAC
GTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGGCCACCATCAC
CATCACCATTAA

Protein sequence (Seq. ID. No. 19)

(Amino-acids corresponding to Prot D fusion partner are in bold)

MDPSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYL
EQDLAMTKDGRLVVIHDHFLDGLTDVAKKFPHRHRKDGRYYVIDFTLK
EIQSLEMTENFETMGGKWSKSSVVGWPTVVRERMRAEPAADGVGAASRDL
EKHGAITSSNTAATNAACAWLEAQEEEEVGFPVTPQVPLRPMTYKAAVDLSH
FLKEKGGLEGLIHSQRRQDILDLYIYHTQGYFPDWQNYTPGPGVRYPLTFGW
CYKLVPVEPDKVEEANKGENTSLLHPVSLHGMDDPEREVLEWRFD SRLAFH
HVARELHPEYFKNCTSGHHHHHHH.

⇒ ProtD-Nef-Tat-HIS

DNA sequence (Seq. ID. No. 20)

THIS PAGE BLANK (USPTO)

Nucleotides corresponding to the Prot D Fusion Partner are in bold.

ATGGATCCAAGCAGCCATTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAA
ATCATTATTGCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATACGTTAGAATCT
AAAGCACTTGCGTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACT
AAGGATGGTCGTTTAGTGGTTATTCACGATCACTTTTATAGATGGCTTGACTGATGTT
GCGAAAAAATCCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTT
ACCTTAAAAGAAATTCAAAGTTTAGAAATGACAGAAAACCTTGAAACCATGGGTGGC
AAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGA
GCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGA
GCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCA
CAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATG
ACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTTAAAAGAAAAGGGGGGACTGGAA
GGGCTAATTCCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACA
CAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTGAGATATCCA
CTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAG
GCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGAT
GACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTTCATCAC
GTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGAGCCAGTAGAT
CCTAGACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAAACTGCTTGTACCAAT
TGCTATTGTAAAAAGTGTTGCTTTCATTGCCAAGTTTGTTCATAACAAAAGCCTTA
GGCATCTCCTATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCTCAAGGCAGT
CAGACTCATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCCGAGGGGACCCG
ACAGGCCCGAAGGAACTAGTGGCCACCATCACCATCACCATTAA

Protein sequence (Seq. ID. No. 21)

(Amino-acids corresponding to Prot D fusion partner are in bold)

MDPSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYLEQDLAMT
KDGRLLVVIHDHFLDGLTDVAKKFPHRHRKDGRYYVIDFTLKEIQSLEMTENFETMGG
KWSKSSVVGWPTVRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAWLEA
QEEEEVGFPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDLWIYHT
QGYFPDWQNYTPGPGVRYPLTFGWICYKLVPVEPDKVEEANKGENTSLLHPVSLHGMD
DPEREVLEWRFD SRLAFHHVARELHPEYFKNCTSEPVDPRLEPWKHPGSQPKTACTN
CYCKKCCFHCQVCFITKALGISYGRKKRRQRRRPPQGSQTHQVSLSKQPTSQSRGDP
TGPKETSGHHHHHH.

⇒ Tat-MUTANT-HIS

DNA sequence (Seq. ID. No. 22)

THIS PAGE BLANK (USPTO)

ATGGAGCCAGTAGATCCTAGACTAGAGCCCTGGAAGCATC 40
 CAGGAAGTCAGCCTAAACTGCTTGCTACCAATTGCTATTG 80
 TAAAAAGTGTGCTTTTCATTGCCAAGTTTGTTCATAACA 120
 GCTGCCTTAGGCATCTCCTATGGCAGGAAGAAGCGGAGAC 160
 AGCGACGAAGACCTCCTCAAGGCAGTCAGACTCATCAAGT 200
 TTCTCTATCAAAGCAACCCACCTCCCAATCCAAAGGGGAG 240
 CCGACAGGCCCCGAAGGAACTAGTGGCCACCATCACCATC 280
 ACCATTAA 288

Protein sequence(Seq. ID. No. 23)

Mutated amino-acids in Tat sequences are in bold.

MEPVDPRLEPWKHPGSQPKTACTNCYCKKCCFHCQVCFIT 40
 AALGISYGRKKRRRQRRRPPQGSQTHQVSLSKQPTSQSKGE 80
 PTGPKETSGHHHHHHH. 95

⇒Nef-Tat-Mutant-HIS

DNA sequence(Seq. ID. No. 24)

ATGGGTGGCAAGTGGTCAAAAAGTAGTGTGGTTGGATGGC 40
 CTACTGTAAGGGAAAGAATGAGACGAGCTGAGCCAGCAGC 80
 AGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACAT 120
 GGAGCAATCACAAGTAGCAATACAGCAGCTACCAATGCTG 160
 CTTGTGCCTGGCTAGAAGCACAAAGAGGAGGAGGTGGG 200
 TTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATGACT 240
 TACAAGGCAGCTGTAGATCTTAGCCACTTTTAAAAGAAA 280
 AGGGGGGACTGGAAGGGCTAATTCCTCCCAACGAAGACA 320
 AGATATCCTTGATCTGTGGATCTACCACACACAAGGCTAC 360
 TTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTCA 400
 GATATCCACTGACCTTTGGATGGTGCTACAAGCTAGTACC 440
 AGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAAGGAGAG 480
 AACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGG 520
 ATGACCCTGAGAGAGAAGTGTAGAGTGGAGGTTTGACAG 560
 CCGCCTAGCATTTTCATCACGTGGCCCGAGAGCTGCATCCG 600
 GAGTACTTCAAGAACTGCACTAGTGAGCCAGTAGATCCTA 640
 GACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAAAC 680
 TGCTTGCTACCAATTGCTATTGTAAGAAAGTGTGCTTTTCAT 720
 TGCCAAGTTTGTTCATAACAGCTGCCTTAGGCATCTCCT 760
 ATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCTCA 800
 AGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCC 840
 ACCTCCCAATCCAAAGGGGAGCCGACAGGCCCCGAAGGAAA 880
 CTAGTGGCCACCATCACCATCACCATTAA 909

THIS PAGE BLANK (USPTO)

Protein sequence (Seq. ID. No. 25)


Mutated amino-acids in Tat sequence are in bold.

MGGKWSKSSVVGWPTVRERMRRAEPAADGVGAASRDLEKH 40
GAITSSNTAATNAACAWLEAQEEEEVGFPVTPQVPLRPMT 80
YKAAVDLSHFLKEKGGLEGLIHSQRRQDILDLWIYHTQGY 120
FPDWQNYTPGPGVRYPLTFGWICYKLVPVEPDKVEEANKGE 160
NTSLLHPVSLHGMDPPEREVLEWRFD SRLAFHHVARELHP 200
EYFKNCTSEPVDPRLEPWKHPGSQPKTACTNCYCKKCCFH 240
CQVCFITAALGISYGRKKRRQRRRPPQGSQTHQVSLSKQP 280
TSQSKGEPTGPKETSGHHHHHH. 302

THIS PAGE BLANK (USPTO)

Figure 2

Map of pRIT14597 integrative vector




MCS POLYLINKER *nef* gene inserted between NcoI and SpeI sites.

Asu II *Nco* I *Spe* I *Eco* RI
 TTCGAA.ACC.ATGGCCGCGGACTAGTGGC.CAC.CAT.CAC.CAT.CAC.CAT.TAA.CGGAATTC
 Thr . Ser . Gly . His . His . His . His . His . His

The amino acid sequence of Figure 3 relates to Seq. ID no. 27 and the nucleic acid sequence of Figure 3 relates to Seq. ID. No. 26.



THIS PAGE BLANK (USPTO)

Figure 3: SDS-PAGE: Nef-Tat-his fusion protein




1: MW (175/83/62,5/47,5/32,5/25/16,5/6,5 kDa)
 2: TNH/23 SP eluate (250 ng)
 3: TNH/23 Purified bulk (250 ng)
 4: TNH/22 Purified bulk (250 ng)

5: MW (175/83/62,5/47,5/32,5/25/16,5/6,5 kDa)
 6: TNH/23 SP eluate (400 ng)
 7: TNH/23 Purified bulk (400 ng)
 8: TNH/22 Purified bulk (400 ng)



THIS PAGE BLANK (USPTO)

Figure 4 : SDS-PAGE: Nef-Tat-his fusion protein



Coomassie blue G250

1: MW (175/83/62,5/47,5/32,5/25/16,5/6,5 kDa)

2: TNH/23 SP eluate (4 μ g)

3: TNH/23 Superdex200 eluate (4 μ g)

4: TNH/23 Purified bulk (4 μ g)


5: TNH/22 Purified bulk (4 μ g)

6: TNH/23 Purified bulk (4 μ g) / non reducing conditions

7: TNH/22 Purified bulk (4 μ g) / non reducing conditions

THIS PAGE BLANK (USPTO)

Figure 6: SDS-PAGE ANALYSIS – reducing conditions
(14% polyacrylamide precasted gels - Novex) See example 5



1: MW (175/83/62/47,5/32,5/25/16,5/6,5 kDa)
2: Purified bulk
3: Purified bulk

THIS PAGE BLANK (USPTO)


Figure 7 (relating to Example 6): SDS-PAGE ANALYSIS:
(4-20% polyacrylamide precasted gels - Novex)

1 2 3 4

5 6 7

1 2 3 4

5 6 7



Coomassie blue G250

Western blot anti Tat

1: MW (175/83/62,5/47,5/32,5/25/16,5/6,5 kDa)

2: Purified bulk (reducing conditions)

3: Purified bulk (reducing conditions)

4: Purified bulk (reducing conditions)


5: Purified bulk (non reducing conditions)

6: Purified bulk (non reducing conditions)

7: Purified bulk (non reducing conditions)

THIS PAGE BLANK (USPTO)

Figure 8 (relating to Example 7): SDS-PAGE ANALYSIS:
(4-20% polyacrylamide precasted gels - Novex)




Coomassie blue G250

Western blot anti Tat


- 1: MW (175/83/62,5/47,5/32,5/25/16,5/6,5 kDa)
- 2: Purified bulk (reducing conditions)
- 3: Purified bulk (non reducing conditions)

THIS PAGE BLANK (USPTO)

FIGURE 9: SDS-PAGE ANALYSIS - REDUCING CONDITIONS
(14% polyacrylamide precasted gels - Novex) see Example 8




Coomassie blue R250




Silver staining

THIS PAGE BLANK (USPTO)

Figure 10: SDS-PAGE ANALYSIS – REDUCING CONDITIONS
(14% polyacrylamide precasted gels - Novex) See Example 9



THIS PAGE BLANK (USPTO)

Figure 11**Map of pRIT14908 integrative vector**

MCS POLYLINKER : NEF gene inserted between NcoI and SpeI sites.

<i>Asu</i> II	<i>Nco</i> I	<i>Spe</i> I	<i>Eco</i> RI
TTCGAA.A	<u>CC.ATG</u>	GCCGCGG	<u>ACTAGT</u>
		.GGC.CAC.CAT.CAC.CAT.CAC.CAT.TAA.CGC	<u>GAATTC</u>
		Thr .Ser . Gly. His . His . His . His . His . His .	

THIS PAGE BLANK (USPTO)

Figure 12

Sequences of Pichia-expressed SIV-NEF-His protein

DNA SEQUENCE:

```

atgggtggagctat t tccatgagggcggtccagggcgtctggagatctgcg      50
acagagactcttg cggggcgcggtggggagacttatgggagactcttaggag      100
aggtggaagatgg atactcgcaatccccaggaggattagacaagggcttg      150
agctcactctctt gtgagggacagaaaatacaatcaggggacagtatatgaa      200
tactccatggaga aacccagctgaagagagagaaaaattagcatacacagaa      250
a caaaaatatgg atgatatagatgaggaagatgatgacttggtaggggta      300
tcagtgaggccaa aagttcccctaagaacaatgagttacaaattggcaat      350
agacatgtctcat tttataaaaagaaaaggggggactggaaagggaatttatt      400
acagtgaagaaga cecatagaatcttagacatatacttagaaaaggaagaa      450
ggcatcataccag attggcaggattacacctcaggaccaggaattagata      500
cccaaagacattt ggctggctatggaaattagtcctgtaaatgtatcag      550
atgaggcacaggag gatgaggagcattattttaatgcatccagctcaaact      600
tcccagtgggatg acccttggggagaggttctagcatggaagtttgatcc      650
aactctggcctac acttatgaggcatatgttagatacccagaagagtttg      700
gaagcaagtcagg ccctgtcagaggaagaggttagaagaaggctaaccgca      750
agaggccttctta acatggctgacaagaaggaaactcgcactagtggcca      800
ccatcaccatcac cattaa.                                         819

```

PROTEIN SEQUENCE:

```


MGGAI SMRRSRPSGDLRQRLLRARGETYGRLLGEVEDGYSQSPGGLDKGL      50
SSLSC EGQKYNQGQYMNTPWRNP AEEREKLAYRKQNMDDIDEEDDDLGVV      100
SVRPKVPLRTMSYKLAIDMSHFIKEKGGLEGIYY SARRHRILDIYLEKEE      150
GIIPDWQDYTS GPGIRYPKTFGWLWKLVPVNV SDEAQEDEEHYLMHPAQT      200
SQWDDPWGEVLAWKFDP TLAYTYEAYVRYPEEFGSKSGLSEEEVRRRLTA      250
RGLLN MADKKETRTSGHHHHHH.                                     272

```

THIS PAGE BLANK (USPTO)

Figure 13

**Coomassie Blue Stained SDS-PAGE of recombinant
Pichia pastoris SIV/NEF expressing strains**



lane 1: P- Y1752 strain

lane 2: S- " "

lane 3: P- Y1772 strain


lane 4: S- " " "

lane 7: P- GS115 strain (negative control)

lane 8: S- " "


THIS PAGE BLANK (USPTO)

Figure 14 (relating to example 13):



THIS PAGE BLANK (USPTO)

Figure 15:



THIS PAGE BLANK (USPTO)